



<https://www.revistaneurocirugia.com>

## O-VAS-21 - Should ruptured aneurysms be treated earlier?

*M. Carvalho, J. Gonçalves, G. Costa, A. Lopes, R. Pereira and M. Barbosa*

*Centro Hospitalar e Universitário de Coimbra.*

### Resumen

**Objectives:** Based in our clinical results and in the recent literature we intend to discuss arguments in favor of an ultra-early treatment policy and a formal protocol proposal.

**Material and methods:** A 12-year database of consecutive ruptured and surgically treated intracranial aneurysms was analysed. Outcome was measured by the Glasgow Outcome Scale. Baseline characteristics were analysed by chi-square and one-way ANOVA tests. The more recent relevant literature was reviewed.

**Results:** 213 patients who were submitted to surgery in the first 72 hours post-ictus were included, 54 of whom have undergone ultra-early surgery. The demographics and pre-operative characteristics of ultra-early and early surgery patients are homogeneous. Overall outcome results are similar for both groups. For the good-grade patients (WFNS grading scale 1 and 2) submitted to ultra-early surgery, a statistically significant ( $p = 0.047$ ) improved 6-month GOS outcome was verified. A trend towards good 6-month GOS outcome was also achieved on Fisher grade 4 patients who were submitted to ultra-early surgery ( $p = 0.052$ ). A low rebleeding rate was observed on the studied population (1.9%). The recent literature on this matter tends to advise treatment of ruptured aneurysms on an emergency basis.

**Conclusions:** Based in our previously reported results and in the literature review a formal protocol for urgent treatment will be discussed bearing in mind local and general restrictions.