



## P186 - EXTRADURAL GIANT HEMANGIOBLASTOMA IN ADOLESCENCE: CASE REPORT AND THERAPEUTIC REVIEW

*J.P. Lavrador, E. Oliveira, W. Teixeira, A. Fernandes Francisco and S. Livraghi*

*Champalimaud Foundation Portugal. Departamento de Neurocirurgia, Hospital Santa Maria, Centro Hospitalar Lisboa Norte.*

### Resumen

**Introduction:** Non-Von Hippel Lindau Spinal Hemangioblastomas usually arise from the vertebral bodies extending to the extradural space and rarely have a purely extradural location, constituting approximately 4% of all spinal extradural tumors.

**Case report:** A 14-year-old male patient was referred to neurosurgical outpatient clinical complaining of raquialgia with no neurological signs. A CT and MRI were done revealing an right extradural intra/extracanal giant hemangioblastoma (22.5 × 11 × 9 cm) extending from D7 to L3 and centered to right D10 foramen with medular compression at this level. The DSA showed an intense tumoural blush and a major feeder exiting at D10 transverse foramen were seen. No embolization was accomplished. A complete resection of extracanal component was made; a D10 right hemilaminectomy was made and an intracanal decompression was achieved; blood loss was minimized due to intratumoral high viscosity and radiopaque cement injection under pressure with radiological control. Post-operative MRI showed a tumoural resection over 95% with a small intracanal remnant and pathology was supportive of hemangioblastoma. The remnant was removed after elective embolization and the patient was returned to his daily living.

**Discussion:** Surgical management of a giant thoracic extradural hemangioblastoma is challenging. A multistep approach is desirable to avoid major blood loss and neurological complications. In terms of surgical approach, cement injection under pressure with radiological control is a safe technique to minimize blood loss.

**Key words:** Spinal hemangioblastoma. Cement injection. Embolization.