



## P160 - STEREOTACTIC CHARACTERIZATION OF ANTERIOR NUCLEUS OF THALAMUS &NDASH; CADAVERIC STUDY

*A. Rainha Campos*<sup>2,3</sup>, *S. Ferreira*<sup>1</sup>, *P. Pereira*<sup>4</sup>, *P. Henriques*<sup>2</sup>, *L. Lucas Neto*<sup>2,5</sup>, *A Gonçalves-Ferreira*<sup>2,3</sup> and *A. Andrade*<sup>1</sup>

<sup>1</sup>Instituto de Biofísica e Engenharia Biomédica, Faculdade de Ciências da Universidade de Lisboa. <sup>2</sup>Instituto de Anatomia, Faculdade de Medicina de Lisboa. <sup>3</sup>Serviço de Neurocirurgia; <sup>4</sup>Laboratório de Neuropatologia, Serviço de Neurologia; <sup>5</sup>Serviço de Neurorradiologia, Hospital de Santa Maria-CHLN, EPE.

### Resumen

**Objectives:** There is a growing interest on the Anterior Nucleus of Thalamus (ANT) as this seems to be a good target for stimulation in refractory limbic epilepsy patients. However, its Magnetic Resonance Imaging (MRI) is not always clear and stereotactic 3 dimensional atlases aren't available. The main purpose of this study was to identify the ANT in a cadaver brain and to build a 3D model in order to calculate its stereotactic coordinates.

**Material and methods:** A single block of a brain containing both thalamus, mammillary bodies and anterior and posterior commissures was studied. The block was cut in a cryomicrotome and the sections were photographed every 500 µm with a hi-resolution camera. After aligning the pictures in MatLab®, a stack of images was created with ImageJ®. GIMP® was used to delineate ANT, mammillary bodies and mammillothalamic tracts of both cerebral hemispheres and Amira® used for segmentation, 3D reconstruction, calculation of volumes and determination of coordinates.

**Results:** Orthogonal views of the block were obtained from 3D reconstructed images with superimposed objects created. Measurements: AC-PC = 24.04 mm; ANT volume = 125.0 mm<sup>3</sup> on the left side and 119.4 mm<sup>3</sup> on the right. Left ANT X = -7.46; Y = 3.49; Z = 11.65; right ANT X = 8.27; Y = 2.73; Z = 13.20.

**Conclusions:** It is possible to create a stereotactic atlas of both hemispheres simultaneously using this method. To further validate and improve accuracy and for extrapolation of data to clinic applications, a brain MRI should be acquired before cutting the block and the study must be replicated with more brains.