




Neurocirugía

<https://www.revistaneurocirugia.com>



C0111 - SYMPTOMATIC ADJACENT SEGMENT DISEASE (ASD) AFTER LUMBAR OR LUMBOSACRAL MINIMALLY INVASIVE INTERBODY FUSION

P. Mendivil¹, J. Lafuente Barraza², D. Manzano², D. Pereira³ and G. Vergara⁴

¹Neurocentro-IMAC Salta, Salta, Argentina. ²Hospital del Mar, Barcelona, Spain. ³Centro Hospitalar Gaia Espinho, Vila Nova de Gaia, Portugal. ⁴Hospital San Bernardo, Salta, Argentina.

Resumen

Objectives: Because of the increase in fusion surgery in recent years, ASD become an important topic in spine surgery. The development of motion preservation technologies and minimally invasive interbody fusion theoretically should lead to a decrease in this pathology.

Methods: We performed a retrospective cohort study based on medical record review. From 2012 to 2016 83 cases of minimally invasive interbody fusion were performed by the authors. Clinical ASD was the principal judgment criterion; the length of follow-up time and initial spinal disease were also recorded.

Results: Only 41 patients met the recruitment requirements or were able to follow (22 M, 19 F) with an average age of 58 (24-78), minimum 6 month follow up, (average 23 month, range 39-6). As preliminary results the incidence of ASD, following MI lumbar fusions in our series was 4.8% (two patients), one with prior adjacent degenerative disease. Probably the incidence will increase with a longer follow up.

Conclusions: Multiple studies have explored the risk factors contributing to ASD that include; Damage to the posterior ligamentous complex, fusion length (especially, three or more levels), sagittal imbalance, high degree of pelvic incidence, facet injury, advanced age, and prior adjacent degenerative disease. Although the limitations of relative short follow up, we found a trend toward decreased risk of ASD compared to the published open group. This suggests that MI LIF may be associated with decreased long-term morbidity regard ASD compared to the open approach. More Data and longer follow up especially after 5 years are need to evaluate the real incidence of ASD after minimally invasive lumbar interbody fusion in comparison with open approach to confirm these findings.