



# Neurocirugía



<https://www.revistaneurocirugia.com>

## P0372 - 30-DAY MORTALITY AND SURVIVAL IN ELDERLY PATIENTS UNDERGOING NEUROSURGERY

*P. Sinha<sup>2</sup>, E. Bligh<sup>2</sup>, D. Smith<sup>3</sup> and Y. Altamimi<sup>3</sup>*

<sup>2</sup>School of Medicine, Sheffield, Yorkshire, UK. <sup>3</sup>Sheffield Teaching Hospitals NHS Trust, Yorkshire, UK.

### Resumen

**Objectives:** The aim of the current study is to investigate 30-day mortality and survival in a cohort of elderly patients following emergency and elective neurosurgery. Design: Retrospective cohort study. Subjects: Patients greater than 70 years admitted to our neurosurgery department between April 2015 and April 2016.

**Methods:** Online patient electronic records were retrieved to gather information. Logistic regression analysis was used to identify predictors of mortality.

**Results:** 388 patients in total of whom 318 were < 80 years (group A) and 70 were > 80 years (group B). Male: Female 1/4 1.2:1. 160 elective and 228 emergency admissions. Overall 30-day mortality = 7.5% (14.3% in group B). Mean survival (95% confidence interval) in group A compared with group B = 497 days (477-516) vs 435 days (383-488) ( $p = 0.014$ ). Overall cumulative survival at one year was 84%. There was a significant difference in cumulative survival at one-year between elective (95%) and emergency (77%) patients. 81% of patients were discharged back to their usual place of residence (this figure was 60% for group B and 70% for emergency patients). Cranial tumour and vascular diagnosis were diagnostic categories that were independent predictors of mortality ( $p < 0.001$  and  $p = 0.01$  respectively).

**Conclusions:** Current selection of elective and emergency elderly patients (including those > 80 years) undergoing neurosurgery is associated with a low 30-day mortality and 1-year survival and a high rate of discharge back to previous residence.