



# Neurocirugía



<https://www.revistaneurocirugia.com>

## P0307 - TRANSCRANIAL CEREBELLAR HERNIATION FOLLOWING CRANIOTOMY

*I. Panero Pérez, C. Eiriz Fernández, D. García Pérez, A. Lagares Abascal, Á. Pérez Nuñez and I. Paredes Sansinenea*

*Hospital 12 de Octubre, Madrid, Spain.*

### Resumen

**Objectives:** To report a case of post-surgical encephalocele through craniotomy burr holes following the resection of a meningioma of the posterior fossa. Postoperative trans-cranial herniation through craniotomy without predisposing conditions it is an uncommon complication.

**Methods:** We report a case of 49-year-old female with history of cephalaea. The CT scan showed a meningioma of the convexity of the posterior fossa. A craniotomy and resection of the lesion was performed without complications. After 30 days the patient consulted in the emergency department referring cephalaea, recurrent vomiting and imbalance. The MRI revealed a trans-cranial cerebellar herniation through craniotomy.

**Results:** An urgent surgery was performed to repair the encephalocele. Post-surgical MRI showed normal position of the posterior fossa structures. At discharge the patient was asymptomatic.

**Conclusions:** Post-surgical encephalocele is an uncommon complication that can occur after the resection of a fossa posterior lesion. To best of our knowledge it is the first case reported with this kind of complication.